§ 520.1846

Administer in molasses block containing 6.6 percent poloxalene, at the rate of 0.8 ounce of block (1.5 grams of poloxalene) per 100 pounds of body weight per day. Provide access to blocks at least 7 days before exposure to bloat-producing conditions.

[40 FR 13838, Mar. 27, 1975, as amended at 40 FR 39857, Aug. 29, 1975; 42 FR 41854, Aug. 19, 1977; 50 FR 5385, Feb. 8, 1985; 54 FR 33501, Aug. 15, 1989; 56 FR 50653, Oct. 8, 1991; 58 FR 26523, May 4, 1993; 60 FR 55659, Nov. 2, 1995]

§ 520.1846 Polyoxyethylene (23) lauryl ether blocks.

- (a) *Specifications.* Each molassesbased block contains 2.2 percent polyoxyethylene (23) lauryl ether.
- (b) *Sponsor*. See No. 050112 in §510.600(c) of this chapter.
- (c) Conditions of use—(1) Amount. 2 grams of polyoxyethylene (23) lauryl ether per 100 kilograms of body weight per day (1 pound of block per 500 kilogram (1,100 pound) animal per day).
- (2) *Indications for use.* For reduction of the incidence of bloat (alfalfa and clover) in pastured cattle.
- (3) Limitations. Administer freechoice to beef cattle and nonlactating dairy cattle only. Initially, provide one block per five head of cattle. Start treatment 10 to 14 days before exposure to bloat-producing pastures. Do not allow cattle access to other sources of salt while being fed this product. Do not feed this product to animals without adequate forage/roughage consumption

 $[50~{\rm FR}~48189,~{\rm Nov.}~22,~1985,~{\rm as~amended~at}~56~{\rm FR}~9841,~{\rm Mar.}~8,~1991]$

§520.1870 Praziquantel tablets.

- (a) Specifications. Each dog tablet contains 34 milligrams (mg) of praziquantel; each cat tablet contains 11.5 or 23 mg of praziquantel.
- (b) *Sponsor*. See No. 000859 in §510.600(c) of this chapter.
- (c) Conditions of use—(1) Dogs—(i) Indications for use. For removal of canine cestodes Dipylidium caninum and Taenia pisiformis. If labeled for use by or on the order of a licensed veterinarian, for removal of the canine cestode Echinococcus granulosus, and for removal and control of the canine cestode Echinococcus multilocularis.

- (ii) Dosage. Dogs 5 pounds and under, ½ tablet (17 mg); 6 to 10 pounds, 1 tablet (34 mg); 11 to 15 pounds, 1½ tablets (51 mg); 16 to 30 pounds, 2 tablets (68 mg); 31 to 45 pounds, 3 tablets (102 mg); 46 to 60 pounds, 4 tablets (136 mg); over 60 pounds, 5 tablets maximum (170 mg).
- (iii) Limitations. Administer directly by mouth or crumbled and in feed. Not intended for use in puppies less than 4 weeks of age. For over-the-counter (OTC) use: Consult your veterinarian before administering tablets to weak or debilitated animals, and for assistance in the diagnosis, treatment, and control of parasitism. For prescription use: Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- (2) Cats—(i) Indications for use. For removal of feline cestodes Dipylidium caninum and Taenia taeniaeformis.
- (ii) *Dosage*. Cats 4 pounds and under, 11.5 mg; 5 to 11 pounds, 23 mg; over 11 pounds, 34.5 mg.
- (iii) *Limitations.* Administer directly by mouth or crumbled and in feed. Not intended for use in kittens less than 6 weeks of age. For OTC use: Consult your veterinarian before administering tablets to weak or debilitated animals, and for assistance in the diagnosis, treatment, and control of parasitism.

[46 FR 60570, Dec. 11, 1981, as amended at 47 FR 26377, June 18, 1982; 55 FR 2234, Jan. 23, 1990; 58 FR 7864, Feb. 10, 1993; 58 FR 42853, Aug. 12, 1993]

§ 520.1871 Praziquantel/pyrantel pamoate tablets.

- (a) Specifications. Each cat tablet contains 18.2 milligrams (mg) praziquantel with 72.6 mg pyrantel (as pyrantel pamoate).
- (b) *Sponsor*. See 000859 in §510.600(c) of this chapter.
- (c) Conditions of use—(1) Cats—(i) Dosage. 1.5 to 1.9 pounds, 1/4 tablet; 2 to 3 pounds, 1/2 tablet; 4 to 8 pounds, 1 tablet; 9 to 12 pounds, 1 1/2 tablets; 13 to 16 pounds, 2 tablets.
- (ii) Indications for use. For removal of tapeworms (Dipylidium caninum and Taenia taeniaeformis), hookworms (Ancylostoma tubaeforme), and large roundworms (Toxocara cati) in cats and kittens.